

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE (DD-MM-YYYY) 6/10/2004		2. REPORT TYPE Final Report		3. DATES COVERED (From - To) 05/01/2000-04/30/03	
4. TITLE AND SUBTITLE Cell-cell interactions, extracellular matrix formation in biofilms and marine fouling				5a. CONTRACT NUMBER NOOO14-00-12-0754	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
				5d. PROJECT NUMBER	
6. AUTHOR(S) Barbara Wigglesworth-Cooksey and Keith E. Cooksey				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Department of Microbiology 109 Lewis Hall Montana State University-Bozeman Bozeman, MT 59717				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Naval Reserach 800 N. Quincy st. Arlington. Va 2217-5000				10. SPONSOR/MONITOR'S ACRONYM(S) ONR	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Distribution unlimited					
<div style="display: flex; justify-content: space-between;"> <div>13. SUPPLEMENTARY NOTES</div> <div style="font-size: 2em; font-weight: bold;">20040617 087</div> </div>					
14. ABSTRACT In the short-term, bacteria isolated from the same marine biofilms as diatoms caused the diatoms to form cellular aggregates, to lose their motility and their ability to adhere to surfaces. Longer incubation times lead to diatom lysis. A result of these actions was the reduction in structural stability of the mixed species biofilm to hydraulic stress. The active compound(s) from the bacteria was found to be secreted into the bacterial growth medium (MW > 10K Daltons) There is sufficient evidence to regard its mode of action as being lectin-like. Experiments with plant-derived fluorescently-conjugated lectins showed that the extracellular polymers of several diatoms stained specifically indicating the presence of galactose, glucose and mannose moieties, but not those of fucose. Our results are significant in terms of the use of bioassays to test the efficacy of candidate antifouling surfaces and studies of littoral sediment stabilization.					
15. SUBJECT TERMS diatoms, bacteria, interactions, biofilms, biofouling, aggregation, confocal microscopy, sediment stability, littoral					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE			Keith Cooksey
unclassified	unclassified	unclassified	unlimited	5	19b. TELEPHONE NUMBER (include area code) 406 994 6136

Cell-Cell interactions, Extracellular Matrix Formation in Biofilms and Marine Fouling

Principal Investigator: Keith E. Cooksey,
Montana State University, Bozeman, MT 59717, phone: (406) 994
6136, fax: (406) 994 4926, email: umbkc@montana.edu

Co-Principal Investigator: Barbara Wigglesworth-Cooksey,
Montana State University, Bozeman, MT 59717
phone: (406) 994 3022, email: 4505barb@montanads1.net

Grant Number: NO0014-00-1-0754

LONG-TERM GOALS

To add to our understanding of the individual and joint roles of bacteria and diatoms in the formation and persistence of marine fouling biofilms and in so doing, provide information that can be used in the design of fouling release coatings. This information concerning the physiological dissection of the biofilm system is generally relevant to other mixed species biofilms, especially those on illuminated marine sediments in the littoral zone.

SPECIFIC OBJECTIVES

To determine in a mixed diatom/bacterial biofilm, which group of organisms influences the production of the extracellular matrix (ECM) that is responsible for biofilm architecture and persistence.

APPROACH

Eukaryote-prokaryote interactions lead to changes in the detection of a surface and potentially none attachment and /or detachment of cells, thus affecting biofilm structure and function. Bacteria and diatoms isolated from the same marine biofilm were used to study this phenomenon. Diatom motility (an analogue of adhesion) and adhesive behavior was monitored as a function of the presence of bacteria and their products. Efforts to quantify the ECM production in single and mixed organism biofilms and to determine its origin were made. Sugar-specific lectins conjugated to fluorophores were used to demonstrate the presence or absence of particular extracellular polymers. Flow-through bioreactors were used to assess biofilm the formation of extracellular products as a function of time using axenic diatom and mixed bacterial/diatom cultures. We also prepared extracellular polymers from diatoms and diatom-colonized surfaces for study with atomic force microscopy by another grantee.

RESULTS AND ACCOMPLISHMENTS

Bioreactor studies showed that diatom biofilms were less

stable in the presence of bacteria than they were in their absence. Specifically *Pseudoalteromonas* sp.4 (16-S RNA measurements indicate a new species) alone, or the medium in which it was grown, caused diatoms to lose motility, adhesion and viability.

As an initial effect, the spent medium caused diatoms (*Amphora coffeaeformis* and *Navicula* sp. 1) to agglutinate (fig1). Extracellular polymers of these diatoms were easily stained with various conjugated lectins, especially those specific for galactose and mannose, but after treatment with bacteria spent medium, the extracellular polymers were not detectable with the lectin.

Navicula sp.1 control and treated




Figure 1. Agglutination effects of *Pseudoalteromonas* sp.4 medium on *Navicula* sp.1

Cells of *Navicula* sp. 1 were treated with spent medium (treated) from *Pseudoalteromonas* sp. 4. The left hand image (untreated) shows typical active dispersal behavior of diatom cells in fresh marine medium. In the right hand image the effect of their re-suspension in medium that had supported bacterial growth can be seen. The 'spent medium' was sterilized by ultra-filtration before the diatoms were re-suspended.

We found that sugars or their polymers (Table 1) protected diatom cells from the effects of the bacterial spent medium. This is strong evidence that the bacterial products capable of agglutinating the diatoms are lectins. A model to explain this phenomenon has been formulated.

Table 1. D-galactose or mannan protects *Navicula* sp.1 from effects of *Pseudoalteromonas* sp. 4

Treatment of diatoms	% motility (n) ¹
(a). ASP ₂ C alone	98 ± 2 (101)
(b). Spent medium from bacteria	42 ± 12 (25) ²
(c). as (b) + 50mM D-galactose	71 ± 7 (46)
(d). as (b) + 7 mg mL ⁻¹ mannan	66 ± 1 (38)

¹After 120 min incubation, (number of cells measured).

Cell density was identical in each experiment, but aggregation of diatoms in the presence of the spent medium of *Pseudoalteromonas* sp. 4 reduced the number of individual motile cells able to be measured. Thus these figures represent the motility of cells not in an aggregate.

² After 180min motility in (b) was reduced to 0% whereas in the other treatments (a, c, and d), motility did not change.

We have shown that lectin staining of diatom extracellular polymers demonstrate differing adhesive structures between species. For instance, whereas *Amphora* [two raphe on the ventral valve] adheres to a surface by two raphe-penetrating pads of polymer, the adhesive pad for *Navicula* sp.1 covers the entire ventral valve surface interface. The adhesive material is obviously secreted from the raphe in *Amphora* but in *Navicula* sp.1 we postulated that it was secreted from valvar pores, i.e., not from the raphe. We have now documented that this is the case.

IMPACT/APPLICATIONS

The use of a single organism type in assessing antifouling coating formulations is not recommended since not all diatom species behave similarly and the influence of indigenous bacteria on the diatom adhesive behavior can be substantial.

HONORS

During the course of this funding period Keith Cooksey has been Chair Elect, Chair and Advisor to section Q of the American Society of Microbiology which deals with Environmental General and Applied Microbiology.

Keith Cooksey has been appointed to Emeritus Status at Montana State University-Bozeman- the first Research Professor at MSU to be so honored.

PUBLICATIONS AND PUBLISHED ABSTRACTS:

Articles

2001

With B. Wigglesworth-Cooksey and D.Berglund, Cell-cell and cell-surface interactions in an illuminated biofilm : Implications for marine sediment stabilization. *Geochemical Trans.*10:e-journal, no pages numbers.

2002.

With B. Wigglesworth-Cooksey, *Diatoms in Biofilms*. Wiley Encyclopedia of Environmental Microbiology, 1051-1063.

2003.

With Arce, F.T., Avci, R., Beech, I.B. and Wigglesworth-Cooksey, B. Microelastic properties of minimally-adhesive surfaces: A comparative study of RTV-11 and Intersleek polymers. *J Phys.Chem*, 119:1671-1682.

With Arce, F.T., Beech, I.B., Avci, R. and Wigglesworth-Cooksey, B. Microelastic domains in minimally-adhesive polymers studied with atomic force microscopy. *Proc.Amer.Soc.Chem.* New Orleans

2004.

With Wigglesworth-Cooksey B. Use of fluorescently-conjugated lectins to study cell-cell interactions in model marine biofilms, Submitted to *Applied and Environmental Microbiology*, April 2004.

With Arce, F.T. Avci, R., Beech I.B. and Wigglesworth-Cooksey, B. A live probe for studying diatom-surface interactions. *Biophysical J.* Submitted May 2004.

Abstracts

2001.

Oxygen tension in the benthic boundary layer : Water clarity, sediment stability and the activity of benthic phototrophs, in "Effects of Hypoxia on Aquatic Biota", [with B.Wigglesworth-Cooksey], January, La Paz, Mexico.

Diatom and bacterial extracellular polymers in sediment stabilization, [With B. Wigglesworth-Cooksey], Division of Biogeochemistry, American Chemical Society, April, San Diego, CA.

2002.

Diatom-bacterial interactions in marine biofilms and their significance in the design of fouling - release coatings, (with B. Wigglesworth-Cooksey). *Annu.Mtg.Amer.Soc.Microbiol.* Salt Lake City, April.

2003.

Bacterial and pennate diatom interactions inhibit diatom adhesive and motile behavior(with B. Wigglesworth-Cooksey), *Annu. Mtg.Amer.Soc.Microbiol.* Washington, DC, May.

Bacterial and pennate diatom interactions via EPS in mixed species biofilms(with B. Wigglesworth-Cooksey), Royal Netherlands Academy for Science and the Arts, Symposium on the

Microphytobenthos, Amsterdam, August.

2004

Involvement of calcium-based signal transduction and fluxes in biofilm formation and cellular dispersal by benthic marine diatoms. (with B. Wigglesworth-Cooksey), Annu.Mtg.Amer.Soc.Microbiol New Orleans, May.